



The Elastic and Radiative Energy Loss Contributions for the Quenching Factor

Luiz Fernando Mackedanz[†]

[†] `luiz.mackedanz@ufrgs.br`

High Energy Phenomenology Group (GFPAE)

Instituto de Física, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

M. B. Gay Ducati, V. P. Gonçalves, L. F. Mackedanz, submitted to Phys. Lett. B

- High- p_{\perp} hadron suppression \rightarrow Jet Quenching;
- Partons propagating through a dense QCD medium created in HIC are expected to suffer energy degradation before their hadronization;
- This energy loss affects the final momentum transverse (p_{\perp}) spectra;
- The quenching factor for parton spectra is expected to be sensitive to the magnitude of the energy loss;
- At RHIC energy regime, the radiative processes really dominate over elastic ones ? What is the critical energy where both mechanisms contribute equally ?



Energy Loss - Mechanisms

- Total energy loss can be decomposed:
 - **Radiative** (gluon bremsstrahlung)
 - ⇒ the particle propagates in the medium and strongly interacts with the background field, radiating gluons;
 - **Collisional** (or elastic)
 - ⇒ the particle experiences multiple elastic scatterings with the partons in the plasma;
- Current works in parton energy loss assume that the radiative energy loss dominates and disregard the collisional one.
- Recently, the inclusion of the elastic energy loss have been considered to account for non-photonic electron data from RHIC.

Radiative Energy Loss



● LCPI formalism (Zakharov)

- describes the evolution of a singlet $q\bar{q}g$ system in a dense medium;
- used to compute the induced gluon emission.

● Induced Gluon Spectrum

$$\frac{dp}{dx} = \int_0^L dz n(z) \frac{d\sigma_{\text{eff}}^{\text{BH}}(x, z)}{dx}.$$

⇒ $n(z)$ → number (partons) density of in the medium (QGP).

● Cross section for gluon production

$$\frac{d\sigma_{\text{eff}}^{\text{BH}}(x, z)}{dx} = \text{Re} \int d\rho \psi^*(\rho, x) \sigma_3(\rho, x) \psi(\rho, x, z),$$

⇒ ρ → transverse distance between quark and gluon;

⇒ σ_3 → cross section for interaction between the $q\bar{q}g$ system and a particle in the medium;



Radiative Energy Loss

GFPA

Light-Cone Wave Functions :

- for transition $q \rightarrow qg$ in the vacuum

$$\psi(\rho, x) = p(x) \left(\frac{\partial}{\partial \rho'_x} - \imath s_g \frac{\partial}{\partial \rho'_y} \right) \int_0^\infty d\xi \exp\left(-\frac{\imath \xi}{L_f}\right) \mathcal{K}_0(\rho, \xi | \rho', 0) |_{\rho'=0}$$

- medium modified, describing transition in the medium

$$\psi(\rho, x, z) = p(x) \left(\frac{\partial}{\partial \rho'_x} - \imath s_g \frac{\partial}{\partial \rho'_y} \right) \int_0^z d\xi \exp\left(-\frac{\imath \xi}{L_f}\right) \mathcal{K}_0(\rho, z | \rho', z - \xi) |_{\rho'=0}$$

$$\Rightarrow p(x) = \imath \sqrt{\alpha_s / 2x} [s_g(2-x) + 2s_q x] / 2M(x);$$

$$\Rightarrow M(x) = Ex(1-x);$$

$$\Rightarrow L_f = 2Ex(1-x)/\epsilon^2, \text{ with } \epsilon^2 = m_g^2(1-x) + m_q^2 x^2$$

$$\Rightarrow s_{q,g} \rightarrow \text{parton helicities}$$

- Green Function for two dimensional Hamiltonian

$$\mathcal{K}_0(\rho_2, z_2 | \rho_1, z_1) = \frac{M(x)}{2\pi \imath (z_2 - z_1)} \exp\left[\frac{\imath M(x)(\rho_2 - \rho_1)^2}{2(z_2 - z_1)}\right],$$

Radiative Energy Loss



Assumptions to evaluate it

- σ_3 can be written in terms of dipole cross section σ_2

$$\sigma_3(\rho, x) = \frac{C_A}{2C_F} [\sigma_2((1-x)\rho) + \sigma_2(\rho) - \frac{1}{C_A^2} \sigma_2(x\rho)] = C_3(x)\rho^2,$$

with

$$C_3(x) = C_2(\rho)A(x)$$

$$A(x) = [1 + (1-x)^2 - \frac{x^2}{N_c^2}] \frac{C_A}{2C_F}$$

- Spectrum is dominated by contributions of the region $\rho \ll 1/\mu$

$$C_2(\rho) \propto \ln\left(\frac{1}{\rho\mu}\right)$$

- Radiative energy loss

$$\Delta E_{\text{rad}} = E \int_{x_{\min}}^{x_{\max}} dx x \frac{dp}{dx}$$

- $x_{\min} = m_g/E$

- $x_{\max} = 1 - m_q/E$



Elastic Energy Loss - conservative view


- First estimated by Bjorken for massless quarks
→ problems in the infrared limit → coherent multiple scatterings
- Problem solved with combination of techniques of classical plasma physics and HTL corrected propagator → Braaten and Thoma
- For heavy quarks (leading order)

$$\frac{dE}{dL} = \frac{8\pi\alpha_s^2 T^2}{3} \left(1 + \frac{n_f}{6}\right) \left[\frac{1}{v} - \frac{1-v^2}{2v^2} \ln\left(\frac{1+v}{1-v}\right) \right] \ln \left[2^{\frac{n_f}{6+n_f}} B(v) \frac{ET}{m_g M} \right] \quad E \ll M$$
$$-\frac{dE}{dL} = \left(1 + \frac{9}{4}\right) \frac{4\pi\alpha_s^2 T^2}{3} \left(1 + \frac{n_f}{6}\right) \ln \left[2^{\frac{n_f}{2(6+n_f)}} 0.92 \frac{\sqrt{ET}}{m_g} \right] \quad E \gg M^2/T$$



- $m_g = \sqrt{\frac{(1+\frac{n_f}{6})}{3}} gT \rightarrow$ thermal gluon mass
- Elastic Energy Loss

$$\Delta E_{\text{elas}} = -c \int_0^L \frac{dE}{dL} dz,$$

$c \Rightarrow$ ad hoc pre factor → used to analyze the effects of different magnitudes




Energy Loss Results - $\alpha_s = 0.5$



⇒ left panel - light quarks - $m_q = 0.1$ GeV

⇒ right panel - charm quarks - $m_q = 1.5$ GeV



Quenching Factor

Basic Idea - particles produced in $A - A$ collisions would have their p_{\perp} -spectra modified with respect to $h - h$ collisions, due this energy loss;

- it is related to the nuclear modification factor $R_{AA}(p_{\perp})$ for hadron spectra;
- transverse momentum of the leading hadron traces the energy of the leading parton;
- high- p_{\perp} partons hadronize at large length scales, outside the medium;
- modifications on the high- p_{\perp} hadron spectra is directly associated to modifications in the parent partons due to energy loss effects;
- Quenching factor

$$\frac{dN^{\text{med}}}{d^2p_{\perp}} = \int d\epsilon D(\epsilon) \frac{dN^{\text{vac}}(p_{\perp} + \epsilon)}{d^2p_{\perp}} \equiv Q(p_{\perp}) \frac{dN^{\text{vac}}(p_{\perp})}{d^2p_{\perp}}$$

- Calculating p_{\perp} distribution in the medium



$$\frac{dN^{\text{med}}}{d^2p_{\perp}} = \frac{1}{2\pi^2 R^2} \int_0^{2\pi} d\phi \int_0^R d^2r \frac{dN^{\text{vac}}(p_{\perp} + \Delta E)}{d^2p_{\perp}}$$

- Geometry

$$L(\phi) = (R^2 - r^2 \sin^2 \phi)^{1/2} - r \cos \phi$$

Quenching factor results

GFI




- Light quarks vacuum parameterization (RHIC hadroproduction data)

$$\frac{dN_L^{\text{vac}}}{d^2p_{\perp}} = A \left(\frac{1}{p_0 + p_{\perp}} \right)^{\nu}$$


- Charm quarks vacuum parameterization (D meson data)

$$\frac{dN_H^{\text{vac}}}{d^2p_{\perp}} = C \left(\frac{1}{bM_c^2 + p_{\perp}^2} \right)^{n/2}$$



Conclusions

- We found that the inclusion of elastic processes can provide corrections to the energy loss calculations;
- For light quarks, the quenching factor is ~ 0.2 , similar to the results for $\pi^0 R_{AA}$ results;
- These results motivate the application for heavy quarks \rightarrow preliminary results indicate larger suppression than predicted by radiative EL calculations;
- Other observables have to be studied to extract the correct magnitude of the contribution of elastic processes to total energy loss;
- Destructive interference between elastic and radiative have been studied in recent works. However, its effect on the elastic energy loss amplitude is still in discussion.



Energy Loss Results - $\alpha_s = 0.4$

