

# Diffractive Higgs boson photoproduction by Double Pomeron Exchange

G.G. Silveira, M.B. Gay Ducati

[gustavo.silveira@ufrgs.br](mailto:gustavo.silveira@ufrgs.br)




**Grupo de Fenomenologia de Partículas de Altas Energias**  
Instituto de Física  
Universidade Federal do Rio Grande do Sul  
Porto Alegre, RS, Brasil

## Motivations

- ▶ The existence of the Higgs boson is an open question in high energy Physics;
- ▶ The Large Hadron Collider will allow to study a new kinematic region never before reached —  $\sqrt{s} = 14 \text{ TeV}$ 
  - ▶ It is expected that the proton-proton collisions in LHC will be able to produce the Higgs boson.
- ▶ Collisions will occur with **no** strong interaction in LHC
  - ▶ The peripheral collisions are a new way to study the Higgs boson production in heavy-ion collisions.
- ▶ Other processes of Higgs production by DPE allow another way to study its production
  - ▶ DPE allows the Higgs boson production by the leading  $ggH$  vertex in the mass range  $M_H = 100 - 200 \text{ GeV}$ .


# Diffractive Higgs production in $pp$ collisions

- ▶ **1991:** Bialas and Landshoff PLB 256 (1991) 540
  - ▶ *Regge Theory* → non-perturbative gluons
- ▶ **1997:** Khoze, Martin and Ryskin PLB 401 (1997) 330
  - ▶ *QCD Pomeron* → two-gluons exchange



# Deeply Virtual Compton Scattering<sup>†</sup>


- ▶ **1998:**  $\gamma^* p \rightarrow \gamma p$  by **Pomeron interaction** in  $ep$  collisions



<sup>†</sup>Frankfurt, Freund, Strikman, PRD **58** (1998) 114001

## Diffractive Higgs photoproduction


- ▶ **Proposal:**  $\gamma p$  process by DPE in  $pp$  collision.



- ▶ The loop is treated in **impact factor formalism** at  $t = 0$ .

## Peripheral collisions


- The  $\gamma p$  process is a subprocess in **peripheral  $pp$  collisions**



- Impact parameter:**  $b > 2R \Rightarrow$  **NO STRONG INTERACTION!**
- Only EM force acts on the second proton  $\Rightarrow$  **REAL PHOTONS**

# Scattering amplitude

- ▶ **Partonic process:**  $\gamma q \rightarrow \gamma + H + q$



- ▶ The scattering amplitude is obtained by the **Cutkosky Rules**

$$\text{Im } \mathcal{A} = \frac{1}{2} \int d(\text{PS})_3 \mathcal{A}_{\text{(left)}} \mathcal{A}_{\text{(right)}} = \frac{20}{9} s \frac{M_H^2 \alpha_s^2 \alpha}{N_c v} \sum_q e_q^2 \underbrace{\left( \frac{\alpha_s C_F}{\pi} \right)}_{*} \int \frac{d\vec{k}^2}{\vec{k}^6}$$

# Cross section for central rapidity

- The cross section is calculated for  $y_H = 0$

$$\frac{d\sigma}{dy_H} \Big|_{y_H=0} = \frac{S_{gap}^2}{18\pi^3 b} \left( \frac{M_H^2}{N_c V} \right)^2 \alpha_s^4 \alpha^2 \left( \sum_q e_q^2 \right)^2 \left[ \int_{Q_0^2}^{\infty} \frac{d\vec{k}^2}{\vec{k}^6} e^{-S(\vec{k}^2, M_H^2)} f_g(x, \vec{k}^2) \right]^2$$


- \* Quark contribution<sup>1</sup>:  $\pi^{-1} \alpha_s C_F \rightarrow f_g(x, \vec{k}^2) = K \partial_{(\ell n \vec{k}^2)} x g(x, \vec{k}^2)$
- Gap Survival Probability<sup>2</sup>:  $S_{gap}^2 \rightarrow 3\% \text{ (5\%)}$  for LHC (Tevatron)
- Gluon radiation suppression<sup>3</sup>: Sudakov factor  $S(\vec{k}^2, M_H^2) \sim \ln^2 \left( M_H^2 / \vec{k}^2 \right)$
- Cutoff  $Q_0^2$ : Necessary to avoid infrared divergencies.
- Electroweak vacuum expectation value:  $v = 246 \text{ GeV}$
- $pT$ -vertex impact parameter:  $b = 5.5 \text{ GeV}^{-2}$

<sup>1</sup>Khoze, Martin, Ryskin, EJPC 14 (2000) 525

<sup>2</sup>Khoze, Martin, Ryskin, EJPC 18 (2000) 167



<sup>3</sup>Forshaw, hep-ph/0508274

## Results: $pp$ vs. $\gamma p$ process




- Greatest event rate in the mass region expected for Higgs detection.

# Results: Gluon distribution functions



- ▶ **Tevatron:**  
Small difference between **LO** and **NLO** distributions;
- ▶ **LHC:**  
NLO distributions show a **greater** contribution than the LO ones.

# Results: Energy dependence



- ▶ **Small** dependence on Higgs mass;
- ▶ Aspect between the **LO** and **NLO** distributions:

▶ **Same** difference in the region  $\underbrace{\sqrt{s} = 9 - 17 \text{ TeV}}_{\text{include LHC region}}$

## Summary

- ▶ Substantial results for Higgs boson production in  $\gamma p$  process present in peripheral collisions;
- ▶ An event rate three times bigger than the rate predicted by previous results in  $p p$  collisions;
- ▶ Clear difference between LO and NLO distributions in the kinematic region of LHC;
- ▶ Necessity of study the cross-section behavior in the very-high-energy limit;
- ▶ Analysis of this production mechanism in  $p p$  collisions:
  - Introduction of photon distribution in the proton;
- ▶ Include QCD and Electroweak theory corrections;
- ▶ Predict physical observables for other production processes.