The physics of AGN-driven galactic outflows

Kastytis Zubovas

kastytis.zubovas@ftmc.lt

Center for Physical Sciences and Technology, Vilnius

FIZINIŲ IR TECHNOLOGIJOS MOKSLŲ CENTRAS Vilnius University Observatory



<ロト <問ト < 目ト < 目

European Week of Astronomy and Space Science, 2017

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	

Outline



2 AGN winds

- Basic wind properties
- Wind shock

3 Large-scale outflows

- Momentum-driven outflow
- Energy-driven outflow
- Wind-outflow connection

Outstanding issues

- Wind mass flow rate
- Temporal AGN variability

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	

Outline



- AGN winds
 - Basic wind properties
 - Wind shock
- 3 Large-scale outflows
 - Momentum-driven outflow
 - Energy-driven outflow
 - Wind-outflow connection
- Outstanding issues
 - Wind mass flow rate
 - Temporal AGN variability

SMBH-galaxy connection ●○○	AGN winds	Large-scale outflows	Outstanding issues	Summary
SMBH-galaxy connection				

ヘロン 人間 とくほど 人ほど 一日

The $M - \sigma$ relation



McConnell & Ma (2013)

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
SMBH-galaxy connection				
The $M - \sigma$ re	lation			



 A connection between SMBH mass and galaxy spheroid stellar velocity dispersion

イロト 不良 とくほ とくほう 二日

McConnell & Ma (2013)

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
SMBH-galaxy connection				

The $M - \sigma$ relation



 A connection between SMBH mass and galaxy spheroid stellar velocity dispersion

• Power law:
$$M \propto \sigma^{\alpha}$$
,
 $\alpha \sim 5$

イロト 不得 トイヨト イヨト 二日

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
SMBH-galaxy connection				

The $M - \sigma$ relation



McConnell & Ma (2013)

- A connection between SMBH mass and galaxy spheroid stellar velocity dispersion
- Power law: $M \propto \sigma^{\alpha}$, $\alpha \sim 5$
- Value of α decreases when galaxies are separated by morphological type

イロト イポト イヨト イヨト

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
SMBH-galaxy connection				

The $M - \sigma$ relation



McConnell & Ma (2013)

- A connection between SMBH mass and galaxy spheroid stellar velocity dispersion
- Power law: $M \propto \sigma^{\alpha}$, $\alpha \sim 5$
- Value of α decreases when galaxies are separated by morphological type
- Implies co-evolution between SMBH and the host galaxy spheroid

イロト イ理ト イヨト イヨト

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	

・ロト ・ 四ト ・ ヨト ・ ヨト

э.

SMBH-galaxy connection

The $M - M_{\rm b}$ relation



Savorgnan et al. (2016)

AGN winds

Large-scale outflows

Outstanding issues

Summary

SMBH-galaxy connection

The $M - M_{\rm b}$ relation



 A connection between SMBH mass and bulge stellar mass

・ロット (雪) (日) (日)

э

Savorgnan et al. (2016)

AGN winds

Large-scale outflows

Outstanding issues

Summary

SMBH-galaxy connection

The $M - M_{\rm b}$ relation



Savorgnan et al. (2016)

- A connection between SMBH mass and bulge stellar mass
- Almost linear (Häring & Rix 2004), at least for ellipticals
- Implies similar growth patterns for SMBH and host galaxies

AGN winds

Large-scale outflows

Outstanding issues

Summary

SMBH-galaxy connection

The $M - M_{\rm b}$ relation



- A connection between SMBH mass and bulge stellar mass
- Almost linear (Häring & Rix 2004), at least for ellipticals
- Implies similar growth patterns for SMBH and host galaxies
- Very large scatter

▲□▶▲□▶▲□▶▲□▶ □ のへで

AGN winds

Large-scale outflows

Outstanding issues

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三■ - のへぐ

Summary

SMBH-galaxy connection

Possible physical SMBH-galaxy connection

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

SMBH-galaxy connection

Possible physical SMBH-galaxy connection

• Connection cannot be gravitational:

$$R_{\rm infl} = \frac{GM_{\rm bh}}{\sigma^2} \simeq 10 M_8 \sigma_{200}^{-2} \text{ pc} \ll R_{\rm b} \simeq 1 \text{ kpc} \qquad (1)$$

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

Summary

SMBH-galaxy connection

Possible physical SMBH-galaxy connection

• Connection cannot be gravitational:

$$R_{\rm infl} = \frac{GM_{\rm bh}}{\sigma^2} \simeq 10 M_8 \sigma_{200}^{-2} \, \rm pc \ll R_b \simeq 1 \, \rm kpc \qquad (1)$$

But can be energetic:

 $E_{\rm accr} \simeq \eta M_{\rm bh} c^2 \simeq 2 \times 10^{61} \eta_{0.1} M_8 \text{ erg} \gg E_{\rm b} \simeq M_{\rm b} \sigma^2 \simeq 8 \times 10^{58} M_{11} \sigma_{200}^2 \text{ erg}$ (2)

AGN winds

Large-scale outflows

Outstanding issues

Summary

SMBH-galaxy connection

Possible physical SMBH-galaxy connection

• Connection cannot be gravitational:

$$R_{\rm infl} = \frac{GM_{\rm bh}}{\sigma^2} \simeq 10 M_8 \sigma_{200}^{-2} \, \rm pc \ll R_b \simeq 1 \, \rm kpc \qquad (1)$$

But can be energetic:

 $E_{\rm accr} \simeq \eta M_{\rm bh} c^2 \simeq 2 \times 10^{61} \eta_{0.1} M_8 \text{ erg} \gg E_{\rm b} \simeq M_{\rm b} \sigma^2 \simeq 8 \times 10^{58} M_{11} \sigma_{200}^2 \text{ erg}$ (2)

• The question then becomes how can energy be communicated to the bulge at a \sim 0.5% efficiency.

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	

Outline

SMBH-galaxy connection

2 AGN winds

- Basic wind properties
- Wind shock

3 Large-scale outflows

- Momentum-driven outflow
- Energy-driven outflow
- Wind-outflow connection

Outstanding issues

- Wind mass flow rate
- Temporal AGN variability

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Basic wind properties				
A 11 11				

Accretion disc wind



 Radiation pressure in the accretion disc drives a wind (King & Pounds 2003):

$$\dot{p}_{w} = \dot{M}_{w} v_{w} = \tau \frac{L_{AGN}}{c}; \quad v_{w} \simeq v_{esc} \simeq \frac{\tau}{\dot{m}} \eta c; \quad \dot{E}_{w} = \frac{\dot{M}_{w} v_{w}^{2}}{2} = \frac{\tau^{2} \eta}{2 \dot{m}} L_{AGN}$$
(3)
$$\eta \equiv \frac{L_{AGN}}{\dot{M}_{AGN} c^{2}}; \quad \dot{m} \equiv \frac{\dot{M}_{w}}{\dot{M}_{acc}}$$
(4)

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@



 Radiation pressure in the accretion disc drives a wind (King & Pounds 2003):

$$\dot{p}_{w} = \dot{M}_{w} v_{w} = \tau \frac{L_{AGN}}{c}; \quad v_{w} \simeq v_{esc} \simeq \frac{\tau}{\dot{m}} \eta c; \quad \dot{E}_{w} = \frac{\dot{M}_{w} v_{w}^{2}}{2} = \frac{\tau^{2} \eta}{2 \dot{m}} L_{AGN}$$
(3)
$$\eta \equiv \frac{L_{AGN}}{\dot{M}_{AGN} c^{2}}; \quad \dot{m} \equiv \frac{\dot{M}_{w}}{\dot{M}_{acc}}$$
(4)

• The wind is quasi-spherical and self-regulates to keep $\tau \simeq 1$ (single-scattering limit) (King 2010)

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@



 Radiation pressure in the accretion disc drives a wind (King & Pounds 2003):

$$\dot{p}_{w} = \dot{M}_{w} v_{w} = \tau \frac{L_{AGN}}{c}; \quad v_{w} \simeq v_{esc} \simeq \frac{\tau}{\dot{m}} \eta c; \quad \dot{E}_{w} = \frac{\dot{M}_{w} v_{w}^{2}}{2} = \frac{\tau^{2} \eta}{2 \dot{m}} L_{AGN}$$
(3)
$$\eta \equiv \frac{L_{AGN}}{\dot{M}_{AGN} c^{2}}; \quad \dot{m} \equiv \frac{\dot{M}_{w}}{\dot{M}_{acc}}$$
(4)

- The wind is quasi-spherical and self-regulates to keep $\tau \simeq 1$ (single-scattering limit) (King 2010)
- Winds likely to be intermittent (e.g. Pounds & Vaughan 2012)

AGN winds

Large-scale outflows

Outstanding issues

▲□▶ ▲□▶ ▲□▶ ▲□▶ = 三 のへで

Summary

Basic wind properties

AGN winds

Large-scale outflows

Outstanding issues

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三■ - のへぐ

Summary

Basic wind properties

Observed winds (UFOs)

• Winds are detected via blueshifted FeXXV and FeXXVI absorption, indicating very high ionisation parameters $\xi \sim 10^6$

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Basic wind properties

- Winds are detected via blueshifted FeXXV and FeXXVI absorption, indicating very high ionisation parameters $\xi \sim 10^6$
- Observed in \sim 40% of local AGN (Tombesi et al. 2010)

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

Summary

Basic wind properties

- Winds are detected via blueshifted FeXXV and FeXXVI absorption, indicating very high ionisation parameters $\xi \sim 10^6$
- Observed in \sim 40% of local AGN (Tombesi et al. 2010)
- Wind velocities fall in range $v_{\rm w} = 0.03 0.3c$ (Reeves et al. 2003, Tombesi et al. 2010)

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

Summary

Basic wind properties

- Winds are detected via blueshifted FeXXV and FeXXVI absorption, indicating very high ionisation parameters $\xi \sim 10^6$
- Observed in \sim 40% of local AGN (Tombesi et al. 2010)
- Wind velocities fall in range $v_{\rm w} = 0.03 0.3c$ (Reeves et al. 2003, Tombesi et al. 2010)
- Spatial extent $r_{\rm w} < 1$ pc

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Wind shock				

▲□▶▲圖▶▲≣▶▲≣▶ ≣ のへの

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Wind shock				

$$T_{\rm sh} = \frac{3\mu m_{\rm p} v_{\rm w}^2}{16k_{\rm B}} \simeq 1.3 \times 10^{10} v_{0.1}^2 \ {\rm K} \tag{5}$$

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ



$$T_{\rm sh} = \frac{3\mu m_{\rm p} v_{\rm w}^2}{16k_{\rm B}} \simeq 1.3 \times 10^{10} v_{0.1}^2 \ {\rm K} \tag{5}$$

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

 Shocked wind cools by inverse Compton scattering against the AGN radiation field, with a cooling time t_C



$$T_{\rm sh} = \frac{3\mu m_{\rm p} v_{\rm w}^2}{16k_{\rm B}} \simeq 1.3 \times 10^{10} v_{0.1}^2 \, {\rm K} \tag{5}$$

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

- Shocked wind cools by inverse Compton scattering against the AGN radiation field, with a cooling time t_C
- Hot wind bubble pushes against the surrounding gas and causes it to expand on a timescale t_{exp}



$$T_{\rm sh} = rac{3\mu m_{
m p} v_{
m w}^2}{16k_{
m B}} \simeq 1.3 imes 10^{10} v_{0.1}^2 \ {
m K}$$
 (5)

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

- Shocked wind cools by inverse Compton scattering against the AGN radiation field, with a cooling time t_C
- Hot wind bubble pushes against the surrounding gas and causes it to expand on a timescale t_{exp}
- $t_{\rm C}/t_{\rm exp}$ increases with radius, determines type of outflow:



$$T_{\rm sh} = rac{3\mu m_{
m p} v_{
m w}^2}{16k_{
m B}} \simeq 1.3 imes 10^{10} v_{0.1}^2 \ {
m K}$$
 (5)

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

- Shocked wind cools by inverse Compton scattering against the AGN radiation field, with a cooling time t_C
- Hot wind bubble pushes against the surrounding gas and causes it to expand on a timescale t_{exp}
- $t_{\rm C}/t_{\rm exp}$ increases with radius, determines type of outflow:
 - Close to the AGN, $t_C/t_{exp} < 1$, cooling is efficient, outflow is driven by wind momentum



$$T_{\rm sh} = rac{3\mu m_{
m p} v_{
m w}^2}{16k_{
m B}} \simeq 1.3 imes 10^{10} v_{0.1}^2 \ {
m K}$$
 (5)

- Shocked wind cools by inverse Compton scattering against the AGN radiation field, with a cooling time t_C
- Hot wind bubble pushes against the surrounding gas and causes it to expand on a timescale t_{exp}
- $t_{\rm C}/t_{\rm exp}$ increases with radius, determines type of outflow:
 - Close to the AGN, $t_C/t_{exp} < 1$, cooling is efficient, outflow is driven by wind momentum
 - Far from the AGN, $t_{\rm C}/t_{\rm exp} > 1$, cooling is inefficient, outflow is driven by wind energy



$$T_{\rm sh} = rac{3\mu m_{
m p} v_{
m w}^2}{16k_{
m B}} \simeq 1.3 imes 10^{10} v_{0.1}^2 \ {
m K}$$
 (5)

- Shocked wind cools by inverse Compton scattering against the AGN radiation field, with a cooling time t_C
- Hot wind bubble pushes against the surrounding gas and causes it to expand on a timescale *t*_{exp}
- $t_{\rm C}/t_{\rm exp}$ increases with radius, determines type of outflow:
 - Close to the AGN, $t_C/t_{exp} < 1$, cooling is efficient, outflow is driven by wind momentum
 - Far from the AGN, $t_C/t_{exp} > 1$, cooling is inefficient, outflow is driven by wind energy
 - The transition radius at which $t_C/t_{exp} = 1$ is known as the cooling radius

AGN winds

Large-scale outflows

Outstanding issues

Summary

Wind shock

AGN winds and outflows



SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Cooling radiu	S			
SMBH-galaxy connection	AGN winds ○○○○●○	Large-scale outflows	Outstanding issues	Summary
------------------------	---------------------	----------------------	--------------------	---------
Wind shock				
Cooling radius				

• If the shocked wind is treated as a uniform plasma, $R_{\rm C} \simeq 500$ pc (King 2003)

SMBH-galaxy connection	AGN winds ○○○○●○	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Cooling radius	6			

- If the shocked wind is treated as a uniform plasma, $R_{\rm C} \simeq 500$ pc (King 2003)
- More realistically, the wind is a two-temperature plasma (Faucher-Giguère & Quataert 2012):

(ロ) (同) (三) (三) (三) (○) (○)

SMBH-galaxy connection	AGN winds ○○○○●○	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Cooling radius	5			

- If the shocked wind is treated as a uniform plasma, $R_{\rm C} \simeq 500$ pc (King 2003)
- More realistically, the wind is a two-temperature plasma (Faucher-Giguère & Quataert 2012):

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

• Electrons cool down very rapidly

SMBH-galaxy connection	AGN winds ○○○○●○	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Cooling radiu	S			

- If the shocked wind is treated as a uniform plasma, $R_{\rm C} \simeq 500$ pc (King 2003)
- More realistically, the wind is a two-temperature plasma (Faucher-Giguère & Quataert 2012):

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

- Electrons cool down very rapidly
- Most of the energy is in protons

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Cooling radiu	2			

- If the shocked wind is treated as a uniform plasma, $R_{\rm C} \simeq 500$ pc (King 2003)
- More realistically, the wind is a two-temperature plasma (Faucher-Giguère & Quataert 2012):
 - Electrons cool down very rapidly
 - Most of the energy is in protons
 - Electron-proton energy exchange inefficient, cooling also inefficient

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Wind shock				
Cooling radiu	IC			

- If the shocked wind is treated as a uniform plasma, $R_{\rm C} \simeq 500$ pc (King 2003)
- More realistically, the wind is a two-temperature plasma (Faucher-Giguère & Quataert 2012):
 - Electrons cool down very rapidly
 - Most of the energy is in protons
 - Electron-proton energy exchange inefficient, cooling also inefficient

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

• Effective *R*_C < 1 pc

AGN winds

Large-scale outflows

Outstanding issues

▲□▶ ▲□▶ ▲□▶ ▲□▶ = 三 のへで

Summary

Wind shock

AGN winds

Large-scale outflows

Outstanding issues

▲□▶ ▲□▶ ▲□▶ ▲□▶ = 三 のへで

Summary

Wind shock

Wind shock in clumpy medium

A realistic ISM has uneven density

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Wind shock

- A realistic ISM has uneven density
- Most of the wind energy escapes through low-density channels (Zubovas & Nayakshin 2014)

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Wind shock

- A realistic ISM has uneven density
- Most of the wind energy escapes through low-density channels (Zubovas & Nayakshin 2014)
- Dense clouds, which can feed the SMBH, are affected mostly by the AGN wind momentum

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

Summary

Wind shock

- A realistic ISM has uneven density
- Most of the wind energy escapes through low-density channels (Zubovas & Nayakshin 2014)
- Dense clouds, which can feed the SMBH, are affected mostly by the AGN wind momentum
- Two types of outflow still occur:

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

Summary

Wind shock

- A realistic ISM has uneven density
- Most of the wind energy escapes through low-density channels (Zubovas & Nayakshin 2014)
- Dense clouds, which can feed the SMBH, are affected mostly by the AGN wind momentum
- Two types of outflow still occur:
 - Dense gas experiences a momentum-driven outflow and evaporation in the surrounding hot diffuse plasma

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Wind shock

- A realistic ISM has uneven density
- Most of the wind energy escapes through low-density channels (Zubovas & Nayakshin 2014)
- Dense clouds, which can feed the SMBH, are affected mostly by the AGN wind momentum
- Two types of outflow still occur:
 - Dense gas experiences a momentum-driven outflow and evaporation in the surrounding hot diffuse plasma
 - Diffuse gas and evaporating dense clouds experience an energy-driven outflow

SMBH-galaxy	connection
000	

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

Summary

Outline

- SMBH-galaxy connection
- 2 AGN winds
 - Basic wind properties
 - Wind shock
- 3 Large-scale outflows
 - Momentum-driven outflow
 - Energy-driven outflow
 - Wind-outflow connection
 - Outstanding issues
 - Wind mass flow rate
 - Temporal AGN variability

AGN winds

Large-scale outflows

Outstanding issues

Summary

AGN winds and outflows



AGN winds

Large-scale outflows

Outstanding issues

Summary

Momentum-driven outflow

Momentum-driven outflow

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─のへで

AGN winds

Large-scale outflows

Outstanding issues

▲□▶ ▲□▶ ▲□▶ ▲□▶ = 三 のへで

Summary

Momentum-driven outflow

Momentum-driven outflow

• Small-scale outflow, $\dot{E} < 10^{-3} L_{\rm AGN}$

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding is
000	000000	• 0 0000	000

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Momentum-driven outflow

Momentum-driven outflow

- Small-scale outflow, $\dot{E} < 10^{-3} L_{AGN}$
- Can only escape to large distances if AGN wind momentum rate is higher than the weight of the dense gas clouds

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Momentum-driven outflow

Momentum-driven outflow

- Small-scale outflow, $\dot{E} < 10^{-3} L_{\rm AGN}$
- Can only escape to large distances if AGN wind momentum rate is higher than the weight of the dense gas clouds
- This condition gives a required luminosity:

$$L_{\rm AGN} > L_{\rm crit} = rac{4 f_{\rm g} c}{G} \sigma^4 \simeq 4.6 imes 10^{46} \sigma_{200}^4 \ {\rm erg \ s^{-1}}$$
 (6)

AGN winds

Large-scale outflows

Outstanding issues

Summary

Momentum-driven outflow

Momentum-driven outflow

- Small-scale outflow, $\dot{E} < 10^{-3} L_{\rm AGN}$
- Can only escape to large distances if AGN wind momentum rate is higher than the weight of the dense gas clouds
- This condition gives a required luminosity:

$$L_{\rm AGN} > L_{\rm crit} = rac{4 f_{\rm g} c}{G} \sigma^4 \simeq 4.6 imes 10^{46} \sigma_{200}^4 \ {\rm erg \ s^{-1}}$$
 (6)

 Assuming that this is the Eddington luminosity, this translates to a critical mass

$$M_{\rm SMBH} > M_{\rm crit} = \frac{f_{\rm g}\kappa}{\pi G^2} \sigma^4 \simeq 3.7 \times 10^8 \sigma_{200}^4 M_\odot \qquad (7)$$

AGN winds

Large-scale outflows

Outstanding issues

Summary

Momentum-driven outflow

Momentum-driven outflow

- Small-scale outflow, $\dot{E} < 10^{-3} L_{AGN}$
- Can only escape to large distances if AGN wind momentum rate is higher than the weight of the dense gas clouds
- This condition gives a required luminosity:

$$L_{\rm AGN} > L_{\rm crit} = rac{4 f_{\rm g} c}{G} \sigma^4 \simeq 4.6 imes 10^{46} \sigma_{200}^4 \ {\rm erg \ s^{-1}}$$
 (6)

Assuming that this is the Eddington luminosity, this translates to a critical mass

$$M_{\rm SMBH} > M_{\rm crit} = \frac{f_{\rm g}\kappa}{\pi G^2} \sigma^4 \simeq 3.7 \times 10^8 \sigma_{200}^4 M_{\odot} \qquad (7)$$

• The calculated $M_{crit}(\sigma)$ is rather similar to, but with a lower exponent value than, the observed $M - \sigma$ correlation

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Momentum-driven outflow				
$M - \sigma$ relation				

▲□▶▲圖▶▲≣▶▲≣▶ ≣ のへの

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Momentum-driven outflow



 Galaxies of different morphological types and in different environments have different ISM and different requirements for the AGN luminosity:

AGN winds

Large-scale outflows

Outstanding issues

Summary

Momentum-driven outflow



- Galaxies of different morphological types and in different environments have different ISM and different requirements for the AGN luminosity:
 - The constant factor in the $M_{\rm crit}(\sigma)$ expression is higher in ellipticals than in spirals and in cluster galaxies than in field galaxies

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

AGN winds

Large-scale outflows

Outstanding issues

Summary

Momentum-driven outflow



- Galaxies of different morphological types and in different environments have different ISM and different requirements for the AGN luminosity:
 - The constant factor in the $M_{\rm crit}(\sigma)$ expression is higher in ellipticals than in spirals and in cluster galaxies than in field galaxies

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

• The same is true for σ

AGN winds

Large-scale outflows

Outstanding issues

Summary

Momentum-driven outflow

$M - \sigma$ relation



Zubovas & King (2013)

- Galaxies of different morphological types and in different environments have different ISM and different requirements for the AGN luminosity:
 - The constant factor in the $M_{\rm crit}(\sigma)$ expression is higher in ellipticals than in spirals and in cluster galaxies than in field galaxies
 - $\bullet~$ The same is true for σ
- The resulting correlation has a steeper slope, closer to observed (Zubovas & King 2013)

・ロット (雪) ・ (日) ・ (日)

ъ

AGN winds

Large-scale outflows

Outstanding issues

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● のへぐ

Summary

Energy-driven outflow

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Energy-driven outflow

Energy-driven outflow

 Once the black hole mass *M*_{SMBH} > *M*_{crit}, or if there is sufficiently diffuse gas close to the SMBH, the outflow becomes large-scale

SMBH-galaxy connection	AGN winds
000	000000

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Energy-driven outflow

- Once the black hole mass *M*_{SMBH} > *M*_{crit}, or if there is sufficiently diffuse gas close to the SMBH, the outflow becomes large-scale
- Shocked wind is adiabatic, $\dot{E}_{\rm out} \simeq \dot{E}_{\rm w} \simeq 0.05 L_{\rm AGN}$

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Energy-driven outflow

- Once the black hole mass *M*_{SMBH} > *M*_{crit}, or if there is sufficiently diffuse gas close to the SMBH, the outflow becomes large-scale
- Shocked wind is adiabatic, $\dot{E}_{\rm out} \simeq \dot{E}_{\rm w} \simeq 0.05 L_{\rm AGN}$
- Typical outflow velocities $v_{out} \sim 10^3$ km s⁻¹, mass flow rates $\dot{M}_{out} \sim 10^3 M_{\odot}$ yr⁻¹

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Energy-driven outflow

- Once the black hole mass *M*_{SMBH} > *M*_{crit}, or if there is sufficiently diffuse gas close to the SMBH, the outflow becomes large-scale
- Shocked wind is adiabatic, $\dot{E}_{\rm out} \simeq \dot{E}_{\rm w} \simeq 0.05 L_{\rm AGN}$
- Typical outflow velocities $v_{out} \sim 10^3$ km s⁻¹, mass flow rates $\dot{M}_{out} \sim 10^3 M_{\odot}$ yr⁻¹
- Outflow momentum $\dot{p}_{\mathrm{out}} \sim 20 L_{\mathrm{AGN}}/c$

AGN winds

Large-scale outflows

Outstanding issues

Summary

Energy-driven outflow

- Once the black hole mass *M*_{SMBH} > *M*_{crit}, or if there is sufficiently diffuse gas close to the SMBH, the outflow becomes large-scale
- Shocked wind is adiabatic, $\dot{E}_{\rm out} \simeq \dot{E}_{\rm w} \simeq 0.05 L_{\rm AGN}$
- Typical outflow velocities $v_{out} \sim 10^3$ km s⁻¹, mass flow rates $\dot{M}_{out} \sim 10^3 M_{\odot}$ yr⁻¹
- Outflow momentum $\dot{p}_{
 m out} \sim 20 L_{
 m AGN}/c$
- These predictions (Zubovas & King 2012) agree very well with observations (Cicone et al. 2014)

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding is:
000	000000	000000	000

Summary

Energy-driven outflow

Observed outflow properties



Cicone et al. (2014)

AGN winds

Large-scale outflows

Outstanding issues

Summary

Energy-driven outflow

Effect on galactic scales

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─のへで

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Energy-driven outflow				

Effect on galactic scales

• Outflow is powerful enough to remove most gas from galaxy spheroid on timescales shorter than star formation timescale (Feruglio et al. 2010, Cicone et al. 2014)

◆□▶ ◆□▶ ▲□▶ ▲□▶ ■ ののの

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	
	000000			

Energy-driven outflow

Effect on galactic scales

- Outflow is powerful enough to remove most gas from galaxy spheroid on timescales shorter than star formation timescale (Feruglio et al. 2010, Cicone et al. 2014)
- Outflowing gas cools rapidly (Zubovas & King 2014, Richings & Faucher-Giguère), t_{cool} <
 t_{dyn}, forms molecular clumps, might lead to star formation within the outflow (Maiolino et al. 2017)

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@
SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summar
000	000000	000000	000	

Energy-driven outflow

Effect on galactic scales

- Outflow is powerful enough to remove most gas from galaxy spheroid on timescales shorter than star formation timescale (Feruglio et al. 2010, Cicone et al. 2014)
- Outflowing gas cools rapidly (Zubovas & King 2014, Richings & Faucher-Giguère), t_{cool} « t_{dyn}, forms molecular clumps, might lead to star formation within the outflow (Maiolino et al. 2017)
- Outflow has very high pressure:

$$\frac{P_{\rm out}}{P_{\rm disc}} \sim \left(\frac{v_{\rm out}}{\sigma}\right)^2 \left(\frac{R_{\rm d}}{R_{\rm out}}\right)^2 \sim 25 \left(\frac{R_{\rm d}}{R_{\rm out}}\right)^2; \quad (8)$$

(ロ) (同) (三) (三) (三) (○) (○)

this compresses gas in the galactic disc and can trigger or enhance star formation there (Zubovas et al. 2013)

AGN winds

Large-scale outflows

Outstanding issues

Summary

Wind-outflow connection

Wind-outflow connection



Tombesi et al. (2015)

SMBH-galaxy	connection
000	

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ◆□▶ ◆□▶ ● ● ● ●

Summary

Outline

- SMBH-galaxy connection
- 2 AGN winds
 - Basic wind properties
 - Wind shock
- 3 Large-scale outflows
 - Momentum-driven outflow
 - Energy-driven outflow
 - Wind-outflow connection
- Outstanding issues
 - Wind mass flow rate
 - Temporal AGN variability

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outsta
000	000000	000000	000

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● のへぐ

Wind mass flow rate

Wind mass flow rate

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Wind mass flow rate				
Wind mass fl	ow rate			

• Model assumption: $\dot{m} \equiv$ 1, i.e. $\dot{M}_{\rm w} \equiv \dot{M}_{\rm acc}$

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● のへぐ

AGN winds

Large-scale outflows

Outstanding issues

Summary

Wind mass flow rate

Wind mass flow rate

- Model assumption: $\dot{m} \equiv 1$, i.e. $\dot{M}_{\rm w} \equiv \dot{M}_{\rm acc}$
- This isn't true in general: disc feeding process doesn't know about the central SMBH mass

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	

Wind mass flow rate

Wind mass flow rate



Naujalis, Zubovas & Semionov (in prep)

- Model assumption: $\dot{m} \equiv 1$, i.e. $\dot{M}_{\rm w} \equiv \dot{M}_{\rm acc}$
- This isn't true in general: disc feeding process doesn't know about the central SMBH mass
- Preliminary 1.5D simulation results suggest that $\dot{m} \sim$ 10; winds have lower $v_{\rm w}$, but higher $\dot{p}_{\rm w}$ and $\dot{E}_{\rm w}$

・ロ ・ ・ 一 ・ ・ 日 ・ ・ 日 ・

3

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summar
000	000000	000000	000	

Wind mass flow rate

Wind mass flow rate



Naujalis, Zubovas & Semionov (in prep)

- Model assumption: $\dot{m} \equiv 1$, i.e. $\dot{M}_{\rm w} \equiv \dot{M}_{\rm acc}$
- This isn't true in general: disc feeding process doesn't know about the central SMBH mass
- Preliminary 1.5D simulation results suggest that $\dot{m} \sim$ 10; winds have lower $v_{\rm w}$, but higher $\dot{p}_{\rm w}$ and $\dot{E}_{\rm w}$
- However, parts of the wind might 'fail' and become the torus, so the escaping wind may have M_w similar to M_{acc}

AGN winds

Large-scale outflows

 $\underset{\circ \bullet \circ}{\text{Outstanding issues}}$

Summary

Temporal AGN variability

Multiple AGN episodes

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─のへで

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Temporal AGN variability

Multiple AGN episodes

 AGN episodes are short (t_{AGN} ~ 10⁵ yr), outflow lifetimes are long (t_{out} > 1 Myr)

AGN winds

Large-scale outflows

Outstanding issues

◆□▶ ◆□▶ ▲□▶ ▲□▶ □ のQ@

Summary

Temporal AGN variability

Multiple AGN episodes

- AGN episodes are short (t_{AGN} ~ 10⁵ yr), outflow lifetimes are long (t_{out} > 1 Myr)
- Outflow correlations might be broken as the AGN fades

AGN winds

Large-scale outflows

Outstanding issues

Summary

Temporal AGN variability

Multiple AGN episodes



- AGN episodes are short (t_{AGN} ~ 10⁵ yr), outflow lifetimes are long (t_{out} > 1 Myr)
- Outflow correlations might be broken as the AGN fades
- Correlations preserved if $L_{
 m AGN}\left(t
 ight) \propto t^{-lpha}$ with exponent $lpha \sim 1$

ヘロト ヘポト ヘヨト ヘヨト

Zubovas (2017, submitted)

AGN winds

Large-scale outflows

Outstanding issues

Summary

Temporal AGN variability

Multiple AGN episodes



- AGN episodes are short (t_{AGN} ~ 10⁵ yr), outflow lifetimes are long (t_{out} > 1 Myr)
- Outflow correlations might be broken as the AGN fades
- Correlations preserved if $L_{
 m AGN}(t) \propto t^{-lpha}$ with exponent $lpha \sim 1$
- Later AGN episodes can illuminate the outflow, but do not break correlations either

Zubovas (2017, submitted)

AGN winds

Large-scale outflows

Outstanding issues

Summary

Temporal AGN variability

Multiple AGN episodes



Zubovas (2017, submitted)

- AGN episodes are short (t_{AGN} ~ 10⁵ yr), outflow lifetimes are long (t_{out} > 1 Myr)
- Outflow correlations might be broken as the AGN fades
- Correlations preserved if $L_{
 m AGN}(t) \propto t^{-lpha}$ with exponent $lpha \sim 1$
- Later AGN episodes can illuminate the outflow, but do not break correlations either
- Observed correlations might be an upper limit to outflow properties

AGN winds

Large-scale outflows

Outstanding issues

Summary

Temporal AGN variability

Correlations as upper limits



SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary

▲□▶▲圖▶▲≣▶▲≣▶ ≣ のQ@

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
Summary				

AGN winds are a powerful source of energy affecting galaxy evolution:

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary



- AGN winds are a powerful source of energy affecting galaxy evolution:
 - AGN wind momentum establishes the *M* σ relation by cutting off the SMBH gas supply

▲□▶ ▲□▶ ▲□▶ ▲□▶ □ のQで

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary

- AGN winds are a powerful source of energy affecting galaxy evolution:
 - AGN wind momentum establishes the $M \sigma$ relation by cutting off the SMBH gas supply
 - Energy-driven AGN outflows clear gas out of galaxies and affect star formation

▲□▶ ▲□▶ ▲三▶ ▲三▶ - 三 - のへで

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	

- AGN winds are a powerful source of energy affecting galaxy evolution:
 - AGN wind momentum establishes the $M \sigma$ relation by cutting off the SMBH gas supply
 - Energy-driven AGN outflows clear gas out of galaxies and affect star formation

◆□▶ ◆□▶ ◆□▶ ◆□▶ ● ● ● ●

 Theoretical predictions agree very well with observed wind and outflow properties

SMBH-galaxy connection	AGN winds	Large-scale outflows	Outstanding issues	Summary
000	000000	000000	000	

- AGN winds are a powerful source of energy affecting galaxy evolution:
 - AGN wind momentum establishes the $M \sigma$ relation by cutting off the SMBH gas supply
 - Energy-driven AGN outflows clear gas out of galaxies and affect star formation
- Theoretical predictions agree very well with observed wind and outflow properties
 - But there are certain complications and outstanding issues

(日) (日) (日) (日) (日) (日) (日)